

**Human RFC2 (a.a.1-354)**


Swiss-Prot #	P35250
Protein names	Replication factor C subunit 2 (Activator 1 subunit 2) (Replication factor C 40 kDa subunit) (RF-C 40 kDa subunit) (RFC40) (Activator 1 40 kDa subunit) (A1 40 kDa subunit)
Gene names	RFC2
Organism	Human

Codon optimized cDNA Clone


cDNA Cat#	RPE-0137
Description	Applications:1. Codon-optimized cDNA sequence designed for recombinant protein production in both E.coli and Mammalian cell. The cDNA has been cloned into E.coli expression vector with 6x Histidine at N-terminus. 2. May be used as RNAi assistance to restore the loss-of-function.
Tag	A 6x histidine tag
Formulation	Liquid
Storage	-20° to -70° C. Avoid repeated freeze-thaw cycles.

Recombinant Protein

Protein Cat#	PE-0137
Source	Recombinant protein derived from E.coli. Or plant cell
Tag	A 6x histidine tag
Formulation	Liquid
Purity	> 90%, as determined by SDS-PAGE under reducing conditions
Activity	N/A
Applications	Western Blot, ELISA or may be used for other applications determined by user
Storage	-20° to -70° C. Avoid repeated freeze-thaw cycles.
Sequence	MEVEAVCGGAGEVEAQSDPAPAFSKAPGSAGHYELPWVEKYRPVKLNEIVGNEDTVSRL EVFAREGNVPNII IAGPPGTGKTTSLCLARALLGPALKDAMLELNASNDRGIDVVRNKI KMFAQQKVTLPKGRHKII IILDEADSMTDGAQQALRRTMETYSKTRFALACNASDKIIEP IQSRCAYLRYTKLTDQILRLMNVIEKERVPTDGLLEAIFTAQGDMRQALNNLQSTF SGFGFINSENVFKVCDEPHLLVKEMIQHCNVANIDEAYKILAHLWHLGYSPEDIIGNIF RVCKTFQMAEYLLKLEFIKEIGYTHMKIAEGVNSLLQAGLLARLCQKTMAPVAS

Magnetic Beads

Fast purifying
DNA, RNA, Peptide
protein, Antibody
[More>>>](#)



Ion Exchange
WCX, SCX, WAX
Affinity Beads
NH₂, COOH, CHO, NHS, I
odoacetyl, SH, OH,
Silica, Thiol,
etc. Conjugate Any
Molecules C₄, C₈,
C₁₈, Endotoxin-
removal, Biotin, Protein
A, G, A/G, L,
Streptavidin,
Monoavidin GST
[More>>>](#)

Thousands
Recombinant
Protein, cDNA
Available>>>



[Place Order >>>](#)

Recombinant-protein-cDNA-Human-Activator-1-subunit-2-RFC40-RFC2

recombinant protein, codon optimized, cDNA, P35250, Human, RFC2, Replication factor, C, subunit, 2, Activator, 1, subunit, 2, Replication factor, C, 40 kDa, subunit, RF-C, 40 kDa, subunit, RFC40, Activator, 1, 40 kDa, subunit, A1, 40 kDa, subunit,

FOR in vitro RESEARCH USE ONLY. NOT FOR USE IN HUMAN